



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The triangle $M_a M_b M_c$ is similar to the triangle ABC .

$$\angle OM_a M_c = \angle QAC.$$

$$\angle OM_b M_c = \angle QBC.$$

$$\angle OM_c M_a = \angle QCA.$$

$\therefore O$ with respect to the triangle $M_a M_b M_c$, is located precisely as Q is with respect to the triangle ABC .

Hence O is Nagel's point of triangle $M_a M_b M_c$.

Also solved by *F. M. McGAW* and *G. B. M. ZERR*.

74. Proposed by **ROBERT J. ALEY, A. M., Ph. D., Professor of Mathematics, Indiana University, Bloomington, Indiana.**

Let O be the center of the inscribed circle. AO produced meets the circumcircle in A' . Find the ratio of AO to OA' .

I. Solution by **WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics, Ohio University, Athens, Ohio.**

The coördinates of A are $\left(\frac{2\Delta}{a}, 0, 0\right)$; of O , (r, r, r) ; and of A' , those of the intersection of $\beta - \gamma = 0 \dots (1)$, with $a\beta\gamma + b\alpha\gamma + c\alpha\beta = 0 \dots (2)$, having the constant relation $a\alpha + b\beta + c\gamma = 2\Delta \dots (3)$. These give for the coördinates

$$\text{of } A' \left(-\frac{(b+c)^2}{a^2} - \frac{2\Delta}{a}, \quad \frac{(b+c)^3}{a^3} + \frac{2\Delta(b+c)}{a^2}, \quad \frac{(b+c)^3}{a^3} + \frac{2\Delta(b+c)}{a^2} \right).$$

The distance d between $(\alpha_1, \beta_1, \gamma_1)$ and $(\alpha_2, \beta_2, \gamma_2)$ is given by

$$d^2 = -\frac{abc}{4\Delta^2} \{a(\beta_1 - \beta_2)(\gamma_1 - \gamma_2) + b(\gamma_1 - \gamma_2)(\alpha_1 - \alpha_2) + c(\alpha_1 - \alpha_2)(\beta_1 - \beta_2)\} \dots (4).$$

Putting $\alpha_1 = (2\Delta/a)$, $\beta_1 = c$, $\gamma_1 = 0$; $\alpha_2 = \beta_2 = \gamma_2 = r$,

$$\overline{AO}^2 = bcr(b+c-a)/2\Delta \dots (5).$$

Putting $\alpha_1, \beta_1, \gamma_1$ equal respectively to the coördinates of A' , and $\alpha_2 = \beta_2 = \gamma_2 = r$ as before, in (4), we get an expression for $\overline{OA'}^2$.

We can then express the ratio of OA to OA' .

II. Solution by **J. SCHEFFER, A. M., Hagerstown, Maryland.**

The point A' is evidently the middle point of arc BC . Since $\angle A'OC = \frac{1}{2}(A+C)$ and $\angle A'CO = \frac{1}{2}(A+B)$, $OA' = A'C = A'B$.

From Ptolemy's theorem, $ACA'B$ being a cyclic quadrilateral,

$$AB \times A'C + AC \times A'B = AA' \times BC, \text{ or}$$

$$c \times OA' + b \times OA' = (AO + OA')a.$$

$$\therefore OA : OA' = b+c-a : a = 3-a : 2a.$$

Also solved by *G. B. M. ZERR* and *CHAS. C. CROSS*.